

AD-A162 574

THE ASYMPTOTIC BEHAVIOR OF AN EXPONENTIAL-TYPE SERIES

1/1

(U) NAVAL WEAPONS CENTER CHINA LAKE CA


W O ALLTOP ET AL OCT 85 NWC-TP-6662 SBI-AD-E900 528

UNCLASSIFIED

F/G 2076

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A162 574

900 528




NWC TP 6662

# The Asymptotic Behavior of an Exponential-Type Series

by  
William O. Altop  
*Research Department*

and  
Charles S. Kenney  
*Information Systems Group*

OCTOBER 1985



NAVAL WEAPONS CENTER  
CHINA LAKE, CA 93550-6001



Approved for public release; distribution is unlimited.

DTIC FILE COPY

85 12 4 041

# Naval Weapons Center

## FOREWORD

This report describes mathematical properties of a function arising in a laser backscattering study. The work was performed at the Naval Weapons Center, China Lake, Calif., during 1985 under Program Element 61152N, Task Area ZR000-01-01, Work Unit 138070.

The report has been reviewed for technical accuracy by D. T. Gillespie.

The authors wish to acknowledge several helpful discussions with H. N. Browne, Jr., D. T. Gillespie, and J. M. Martin of NWC.

Approved by  
E. B. ROYCE, *Head*  
Research Department  
23 September 1985

Under authority of  
K. A. DICKERSON  
Capt., USN  
*Commander*

Released for publication by  
B. W. HAYS  
*Technical Director*

NWC Technical Publication 6662

Published by	Technical Information Department
Collation	Cover, 4 leaves
First printing	30 copies

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b RESTRICTIVE MARKINGS		
2a SECURITY CLASSIFICATION AUTHORITY			3 DISTRIBUTION / AVAILABILITY OF REPORT A Statement: approved for public release; distribution is unlimited.		
2b DECLASSIFICATION / DOWNGRADING SCHEDULE					
4 PERFORMING ORGANIZATION REPORT NUMBER(S) NWC TP 6662			5 MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Naval Weapons Center		6b OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c ADDRESS (City, State, and ZIP Code) China Lake, CA 93555-6001			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Naval Weapons Center		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c ADDRESS (City, State, and ZIP Code) China Lake, CA 93555-6001			10 SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO 61152N	PROJECT NO	TASK NO ZR000-01-01
			WORK UNIT NO 138070		
11 TITLE (Include Security Classification) THE ASYMPTOTIC BEHAVIOR OF AN EXPONENTIAL-TYPE SERIES					
12 PERSONAL AUTHOR(S) Alltop, William O. and Charles S. Kenney					
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM 85 Jan TO 85 June		14. DATE OF REPORT (Year, Month, Day) 1985, September	
15 PAGE COUNT 6					
16 SUPPLEMENTARY NOTATION					
17 COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
12	01		Laser Backscatter, Convex Functions, Exponential Series		
19 ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>(U) The limit of a certain function of three variables is shown to be exponential in one variable and independent of the other two. The function is a generalization of an infinite series arising in the calculation of backscattering of sharp laser pulses in an infinite cloud of isotropic scatterers. Convexity of the function in a particular region enables the evaluation of the asymptotic behavior without extensive algebraic manipulations.</p>					
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a NAME OF RESPONSIBLE INDIVIDUAL William O. Alltop			22b TELEPHONE (Include Area Code) 619-939-3034		22c OFFICE SYMBOL NWC Code 3807

## 1. INTRODUCTION

The purpose of this report is to show that

$$\lim_{x \rightarrow \infty} \left[ e^{-x} E(x, \alpha) \right] = 1 \quad \text{for } -1 \leq \alpha \leq 1 \quad (1)$$

where

$$E(x, \alpha) \equiv \sum_{k=0}^{\infty} \left( \frac{k+1}{x} \right)^{\alpha} \frac{x^k}{k!} \quad \text{for } x > 0 \text{ and all } \alpha$$

More specifically, we will show that

$$e^x + \alpha \leq E(x, \alpha) \leq e^x + \frac{\alpha}{x} e^x \quad \text{for } 0 \leq \alpha \leq 1 \quad (2)$$

and

$$e^x + \frac{\alpha}{x} e^x \leq E(x, \alpha) \leq e^x + \alpha \quad \text{for } -1 \leq \alpha \leq 0 \quad (3)$$

As a corollary to Equation 1 we have

$$\lim_{x \rightarrow \infty} \left[ x^{-1/2} e^{-x} S(x) \right] = 1 \quad (4)$$

where

$$S(x) \equiv \sum_{k=0}^{\infty} (k+1)^{1/2} \frac{x^k}{k!} = x^{1/2} E(x, \frac{1}{2})$$

Indeed it was an investigation of the asymptotic behavior of  $S(x)$  that led to this study.

The series  $S(x)$  arose in a calculation of time-dependent backscattering of sharp laser pulses in an infinite cloud of isotropic scatterers (Reference 1). More specifically, letting  $x$  denote the time (suitably nondimensionalized) after pulse emission, it turns out that the intensity of the total backscattered signal at times  $x \gg 1$  is approximately proportional to  $x^{-1/2} e^{-\alpha} S(x)$  where  $\alpha \geq 1$ . A sensible interpretation of this result evidently requires a knowledge of the behavior of  $S(x)$  for large  $x$ . The limit (Equation 4) implies that the backscattered intensity at times  $x \gg 1$  is proportional to  $x^{-3/2} e^{-(\alpha-1)x}$ .

In Section 2 we prove Equations 2 and 3 by considering  $E(x, \alpha)$  as a function of  $\alpha$  for fixed  $x > 0$ . This turns out to be a convex function which is easily evaluated at  $\alpha = -1, 0$  and  $1$ . These values together with the convexity property yield Equations 2 and 3.

In Section 3 we prove the slightly improved inequality

$$E(x, \frac{1}{2}) \leq (1 + x^{-1})^{\frac{1}{2}} e^{\frac{x}{2}}$$

This results from manipulating the series for  $(E(x, \frac{1}{2}))^2$ .

In Section 4  $E(x, \alpha)$  is generalized by the addition of one parameter, and the corresponding generalizations of Equations 1 through 3 are proved. Throughout it will be assumed that  $x > 0$ .

## 2. BOUNDS FOR $E(x, \alpha)$

For all  $\alpha$  and fixed  $x > 0$ , the series representation for  $E(x, \alpha)$  can be differentiated term by term with respect to  $\alpha$  due to uniform convergence.

$$E'(x, \alpha) = \sum_{k=0}^{\infty} \left( \frac{k+1}{x} \right)^{\alpha} \ln \left( \frac{k+1}{x} \right) \frac{x^k}{k!}$$

$$E''(x, \alpha) = \sum_{k=0}^{\infty} \left( \frac{k+1}{x} \right)^{\alpha} \left[ \ln \left( \frac{k+1}{x} \right) \right]^2 \frac{x^k}{k!}$$

$E(x, \alpha)$  is a convex function of  $\alpha$  since  $E''(x, \alpha) > 0$  for all  $\alpha$ . It is easy to show that  $E(x, -1) = e^{-x} - 1$  and  $E(x, 0) = e^{-x}$ . The following steps show the less obvious evaluation of  $E(x, 1)$ :

$$\begin{aligned} E(x, 1) &= \sum_{k=0}^{\infty} \left( \frac{k+1}{x} \right) \frac{x^k}{k!} \\ &= \frac{1}{x} + \sum_{k=1}^{\infty} \left( 1 + \frac{1}{k} \right) \frac{x^{k-1}}{(k-1)!} \\ &= \frac{1}{x} + \sum_{k=1}^{\infty} \frac{x^{k-1}}{(k-1)!} + \frac{1}{x} \sum_{k=1}^{\infty} \frac{x^k}{k!} \\ &= \frac{1}{x} + e^x + \frac{1}{x} (e^x - 1) \end{aligned}$$

$$= e^x + \frac{1}{x}e^x$$

For fixed  $x$  we have points  $P(-1)$ ,  $P(0)$ , and  $P(1)$  on the curve  $E = E(x, \alpha)$  in the  $\alpha E$ -plane:

$$P(-1) = (-1, e' - 1)$$

$$P(0) = (0, e')$$

$$P(1) = (1, e' + e'/x)$$

Letting  $L_k$  denote the line joining  $P(k-1)$  and  $P(k)$  for  $k=0, 1$ , we obtain the following equations for  $L_k$ :

$$L_0: E = e' + \alpha$$

$$L_1: E = e' + \alpha e'/x$$

Since  $E(x, \alpha)$  is convex and the three points  $P(k)$  lie on the curve  $E = E(x, \alpha)$ , the curve must lie between  $L_0$  and  $L_1$  in the interval  $-1 \leq \alpha \leq 1$ . This proves Equations 2 and 3.

### 3. AN IMPROVED UPPER BOUND

For the case  $\alpha = \frac{1}{2}$ , the upper bound in Equation 2 can be slightly improved by computing  $(E(x, \frac{1}{2}))^2$ .

$$\begin{aligned} (E(x, \frac{1}{2}))^2 &= \sum_{k=0}^{\infty} \sum_{r=0}^k \left( \frac{r+1}{x} \right)^{\frac{1}{2}} \frac{x^r}{r!} \left( \frac{k-r+1}{x} \right)^{\frac{1}{2}} \frac{x^{k-r}}{(k-r)!} \\ &= \sum_{k=0}^{\infty} \frac{x^{k-1}}{k!} \sum_{r=0}^k \binom{k}{r} [(r+1)(k-r+1)]^{\frac{1}{2}} \end{aligned}$$

For  $0 \leq r \leq k$ ,

$$(r+1)(k-r+1) \leq \left( \frac{k+2}{2} \right)^2$$

with equality holding for  $r = k/2$ . Therefore,

Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	





$$\begin{aligned}
\left(E(x, \tfrac{1}{2})\right)^2 &< \sum_{k=0}^{\infty} \frac{x^{k-1}}{k!} \left(\frac{k+2}{2}\right) \sum_{r=0}^k \binom{k}{r} \\
&= \sum_{k=0}^{\infty} \frac{x^{k-1}}{k!} \left(\frac{k+2}{2}\right) 2^k = \sum_{k=0}^{\infty} \frac{(2x)^{k-1}}{k!} (k+2) \\
&= \frac{1}{x} + \sum_{k=1}^{\infty} \frac{(2x)^{k-1}}{(k-1)!} + \frac{1}{x} \sum_{k=1}^{\infty} \frac{(2x)^k}{k!} \\
&= \frac{1}{x} + e^{2x} + \frac{1}{x} (e^{2x} - 1) = \left(1 + \frac{1}{x}\right) e^{2x}
\end{aligned}$$

It follows that

$$E(x, \tfrac{1}{2}) \leq \left(1 + \frac{1}{x}\right)^{\frac{1}{2}} e^x < \left(1 + \frac{1}{2x}\right) e^x$$

#### 4. A GENERALIZATION

Let  $E_A(x, \alpha)$  be defined by

$$E_A(x, \alpha) = \sum_{k=0}^{\infty} \left(\frac{k+A}{x}\right)^{\alpha} \frac{x^k}{k!}$$

for  $A \geq 1$  and all  $\alpha$ , so that  $E(x, \alpha) = E_1(x, \alpha)$ .

For fixed  $A$  and  $x$ ,  $E_A(x, \alpha)$  is a convex function of  $\alpha$ , assuming the following values at  $\alpha = -1$ , 0, and 1:

$$E_A(x, -1) = e^x - F_A(x)$$

$$E_A(x, 0) = e^x$$

$$E_A(x, 1) = e^x + A e^x / x$$

where

$$F_A(x) = (A-1) \sum_{k=0}^{\infty} \left(\frac{1}{k+A-1}\right) \frac{x^k}{k!}$$

Defining the three points  $P(-1)$ ,  $P(0)$ ,  $P(1)$  and the lines  $L_{-1}$  and  $L_1$  as in Section 2, we have the following equations:

$$L_0: E = e^x + \alpha F_A(x)$$

$$L_1: E = e^x + \alpha A e^x/x$$

The curve  $E = E_A(x, \alpha)$  is bounded by  $L_0$  and  $L_1$  in the  $\alpha E$ -plane. It follows, as in Section 2, that

$$e^x + \alpha F_A(x) \leq E_A(x, \alpha) \leq e^x + \frac{\alpha A}{x} e^x \quad \text{for } 0 \leq \alpha \leq 1$$

$$e^x + \frac{\alpha A}{x} e^x \leq E_A(x, \alpha) \leq e^x + \alpha F_A(x) \quad \text{for } -1 \leq \alpha \leq 0$$

and

$$\lim_{x \rightarrow \infty} \left| e^{-x} E_A(x, \alpha) \right| = 1 \quad \text{for } -1 \leq \alpha \leq 1 \text{ and } A \geq 1$$

We also obtain the following bound for  $F_A(x)$ :

$$F_A(x) \leq \frac{A}{x} e^x \quad \text{for } A \geq 1$$

REFERENCE

1. D. T. Gillespie. "A Calculation of n-Scattered Lidar Returns for Large n in an Idealized Cloud,"  
*J. Opt. Soc. Am. A.* (to be published).

**END**

**FILMED**

---

*1-86*

**DTIC**